

Borderline Personality Disorder

Samyak Makwana
R R Lalan College, Bhuj

Abstract

Personality Disorder is very difficult to diagnose. Also it is very difficult to manage. Borderline personality disorder is one type of personality disorder which is seen in most of the people. In this disorder a person can not regulate his emotions, impulsiveness and unstable relationship with others. Therefore in the therapeutic session psychologist role is to eliminate these three symptoms or to decrease this symptoms. Sometimes person with Borderline Personality Disorder try to selfharm. Dialectical Behavior Therapy implies to treat the Borderline Personality Disorder. There are clusters of symptoms in this disorder. It takes long period to manage these symptoms. This article presents the definition, symptoms and treatments of Borderline Personality Disorder.

Keywords: Borderline Personality Disorder, Self harm, Impulsiveness, Dialectical Behavior Therapy

Borderline personality disorder is a serious mental illness marked by unstable moods, behavior, and relationships. In 1980, the *Diagnostic and Statistical Manual for Mental Disorders, Third Edition* (DSM-III) listed borderline personality disorder as a diagnosable illness for the first time. Most psychiatrists and other mental health professionals use the DSM to diagnose mental illnesses. Because some people with severe borderline personality disorder have brief psychotic episodes, experts originally thought of this illness as atypical, or borderline, versions of other mental disorders.¹ While mental health experts now generally agree that the name “borderline personality disorder” is misleading, a more accurate term does not exist yet.

Most people who have borderline personality disorder suffer from:

- Problems with regulating emotions and thoughts
- Impulsive and reckless behavior
- Unstable relationships with other people.

People with this disorder also have high rates of co-occurring disorders, such as depression, anxiety disorders, substance abuse, and eating disorders, along with self-harm, suicidal behaviors, and completed suicides. According to data from a subsample of participants in a national survey on mental disorders, about 1.6 percent of adults in the United States have borderline personality disorder in a given year.² Borderline personality disorder is often viewed as difficult to treat. However, recent research shows that borderline personality disorder can be treated effectively, and that many people with this illness improve over time.^{1, 3, 4}

What are the symptoms of borderline personality disorder?

According to the DSM, Fourth Edition, Text Revision (DSM-IV-TR), to be diagnosed with borderline personality disorder, a person must show an enduring pattern of behavior that includes at least five of the following symptoms:

Extreme reactions—including panic, depression, rage, or frantic actions—to abandonment, whether real or perceived

A pattern of intense and stormy relationships with family, friends, and loved ones, often veering from extreme closeness and love (idealization) to extreme dislike or anger (devaluation)

Distorted and unstable self-image or sense of self, which can result in sudden changes in feelings, opinions, values, or plans and goals for the future (such as school or career choices)

Impulsive and often dangerous behaviors, such as spending sprees, unsafe sex, substance abuse, reckless driving, and binge eating

Recurring suicidal behaviors or threats or self-harming behavior, such as cutting

Intense and highly changeable moods, with each episode lasting from a few hours to a few days

Chronic feelings of emptiness and/or boredom

Inappropriate, intense anger or problems controlling anger

Having stress-related paranoid thoughts or severe dissociative symptoms, such as feeling cut off from oneself, observing oneself from outside the body, or losing touch with reality.

Suicide and Self-harm

Self-injurious behavior includes suicide and suicide attempts, as well as self-harming behaviors, described below. As many as 80 percent of people with borderline personality disorder have suicidal behaviors,⁷ and about 4 to 9 percent commit suicide.^{4,7} Suicide is one of the most tragic outcomes of any mental illness. Some treatments can help reduce suicidal behaviors in people with borderline personality disorder. For example, one study showed that dialectical behavior therapy (DBT) reduced suicide attempts in women by half compared with other types of psychotherapy, or talk therapy. DBT also reduced use of emergency room and inpatient services and retained more participants in therapy, compared to other approaches to treatment.⁷ For more information about DBT, see the section, “How is borderline personality disorder treated?” Unlike suicide attempts, self-harming behaviors do not stem from a desire to die. However, some self-harming behaviors may be life threatening. Self-harming behaviors linked with borderline personality disorder include cutting, burning, hitting, head banging, hair pulling, and other harmful acts. People with borderline personality disorder may self-harm to help regulate their emotions, to punish themselves, or to express their pain.⁸ They do not always see these behaviors as harmful.

When does borderline personality disorder start?

Borderline personality disorder usually begins during adolescence or early adulthood.^{1, 9} Some studies suggest that early symptoms of the illness may occur during childhood.^{10, 11} Studies suggest early symptoms may occur in childhood. Some people with borderline personality disorder experience severe symptoms and require intensive, often inpatient, care. Others may use some outpatient treatments but never need hospitalization or emergency care. Some people who develop this disorder may improve without any treatment.¹²

What illnesses often co-exist with borderline personality disorder?

Borderline personality disorder often occurs with other illnesses. These co-occurring disorders can make it harder to diagnose and treat borderline personality disorder, especially if symptoms of other illnesses overlap with the symptoms of borderline personality disorder. Women with borderline personality disorder are more likely to have co-occurring disorders such as major depression, anxiety disorders, or eating disorders. In men, borderline personality disorder is more likely to co-occur with disorders such as substance abuse or antisocial personality disorder.¹³ According to the NIMH-funded National Comorbidity Survey Replication—the largest national study to date of mental disorders in U.S. adults—about 85 percent of people with borderline personality disorder also meet the diagnostic criteria for another mental illness.² Other illnesses that often occur with BPD include diabetes, high blood pressure, chronic back pain, arthritis, and fibromyalgia.^{14,}

¹⁵ These conditions are associated with obesity, which is a common side effect of the medications prescribed to treat borderline personality disorder and other mental disorders. For more information, see the section, “How is borderline personality disorder treated?”

What are the risk factors for borderline personality disorder?

Research on the possible causes and risk factors for borderline personality disorder is still at a very early stage. However, scientists generally agree that genetic and environmental factors are likely to be involved. Studies on twins with borderline personality disorder suggest that the illness is strongly inherited.^{16, 17} Another study shows that a person can inherit his or her temperament and specific personality traits, particularly impulsiveness and aggression.¹⁸ Scientists are studying genes that help regulate emotions and impulse control for possible links to the disorder.¹⁹ Social or cultural factors may increase the risk for borderline personality disorder. For example, being part of a community or culture in which unstable family relationships are common may increase a person’s risk for the disorder.¹ Impulsiveness, poor judgment in lifestyle choices, and other consequences of BPD may lead individuals to risky situations. Adults with borderline personality disorder are considerably more likely to be the victim of violence, including rape and other crimes.

How is borderline personality disorder diagnosed?

Unfortunately, borderline personality disorder is often underdiagnosed or misdiagnosed.^{20, 21} A mental health professional experienced in diagnosing and treating mental disorders—such as a psychiatrist, psychologist, clinical social worker, or psychiatric nurse—can detect borderline personality disorder based on a thorough interview and a discussion about symptoms. A careful and thorough medical exam can help rule out other possible causes of symptoms. The mental health professional may ask about symptoms and personal and family medical histories, including any history of mental illnesses. This information can help the mental health professional decide on the best treatment. In some cases, co-occurring mental illnesses may have symptoms that overlap with borderline personality disorder, making it difficult to distinguish

borderline personality disorder from other mental illnesses. For example, a person may describe feelings of depression but may not bring other symptoms to the mental health professional's attention.

No single test can diagnose borderline personality disorder. Scientists funded by NIMH are looking for ways to improve diagnosis of this disorder. One study found that adults with borderline personality disorder showed excessive emotional reactions when looking at words with unpleasant meanings, compared with healthy people. People with more severe borderline personality disorder showed a more intense emotional response than people who had less severe borderline personality disorder.⁶

What studies are being done to improve the diagnosis of borderline personality disorder?

Recent neuroimaging studies show differences in brain structure and function between people with borderline personality disorder and people who do not have this illness.^{22, 23} Some research suggests that brain areas involved in emotional responses become overactive in people with borderline personality disorder when they perform tasks that they perceive as negative.²⁴ People with the disorder also show less activity in areas of the brain that help control emotions and aggressive impulses and allow people to understand the context of a situation. These findings may help explain the unstable and sometimes explosive moods characteristic of borderline personality disorder.^{19, 25} Another study showed that, when looking at emotionally negative pictures, people with borderline personality disorder used different areas of the brain than people without the disorder. Those with the illness tended to use brain areas related to reflexive actions and alertness, which may explain the tendency to act impulsively on emotional cues.²⁶ These findings could inform efforts to develop more specific tests to diagnose borderline personality disorder.⁶

How is borderline personality disorder treated?

Borderline personality disorder can be treated with psychotherapy, or "talk" therapy. In some cases, a mental health professional may also recommend medications to treat specific symptoms. When a person is under more than one professional's care, it is essential for the professionals to coordinate with one another on the treatment plan. The treatments described below are just some of the options that may be available to a person with borderline personality disorder. However, the research on treatments is still in very early stages. More studies are needed to determine the effectiveness of these treatments, who may benefit the most, and how best to deliver treatments.

Psychotherapy

Psychotherapy is usually the first treatment for people with borderline personality disorder. Current research suggests psychotherapy can relieve some symptoms, but further studies are needed to better understand how well psychotherapy works.²⁷ It is important that people in therapy get along with and trust their therapist. The very nature of borderline personality disorder can make it difficult for people with this disorder to maintain this type of bond with their therapist.²⁸

Types of psychotherapy used to treat borderline personality disorder include the following:

1. Cognitive behavioral therapy (CBT). CBT can help people with borderline personality disorder identify and change core beliefs and/or behaviors that underlie inaccurate perceptions of themselves and others and problems interacting with others. CBT may help reduce a range of mood and anxiety symptoms and reduce the number of suicidal or self-harming behaviors.²⁹

2. Dialectical behavior therapy (DBT). This type of therapy focuses on the concept of mindfulness, or being aware of and attentive to the current situation. DBT teaches skills to control intense emotions, reduces self-destructive behaviors, and improves relationships. This therapy differs from CBT in that it seeks a balance between changing and accepting beliefs and behaviors.³⁰

3. Schema-focused therapy. This type of therapy combines elements of CBT with other forms of psychotherapy that focus on reframing schemas, or the ways people view themselves. This approach is based on the idea that borderline personality disorder stems from a dysfunctional self-image—possibly brought on by negative childhood experiences—that affects how people react to their environment, interact with others, and cope with problems or stress.³¹

Therapy can be provided one-on-one between the therapist and the patient or in a group setting. Therapist-led group sessions may help teach people with borderline personality disorder how to interact with others and how to express themselves effectively. One type of group therapy, Systems Training for Emotional Predictability and Problem Solving (STEPPS), is designed as a relatively brief treatment consisting of 20 two-hour sessions led by an experienced social worker. Scientists funded by NIMH reported that STEPPS, when used with other types of treatment (medications or individual psychotherapy), can help reduce symptoms and problem behaviors of borderline personality disorder, relieve symptoms of depression, and improve quality of life.³² The effectiveness of this type of therapy has not been extensively studied. Families of people with borderline personality disorder may also benefit from therapy. The challenges of dealing with an ill relative on a daily basis can be very stressful, and family members may unknowingly act in ways that worsen their relative's symptoms.

Some therapies, such as DBT-family skills training (DBTFST), include family members in treatment sessions. These types of programs help families develop skills to better understand and support a relative with borderline personality disorder. Other therapies, such as Family Connections, focus on the needs of family members. More research is needed to determine the effectiveness of family therapy in borderline personality disorder. Studies with other mental disorders suggest that including family members can help in a person's treatment.³³ Other types of therapy not listed in this booklet may be helpful for some people with borderline personality disorder. Therapists often adapt psychotherapy to better meet a person's needs. Therapists may switch from one type of therapy to another, mix techniques from different therapies, or use a combination therapy. Some symptoms of borderline personality disorder may come and go, but the core symptoms of highly changeable moods, intense anger, and impulsiveness tend to be more persistent.³⁴ People whose symptoms improve may continue to face issues related to co-occurring disorders, such as depression or post-traumatic stress disorder.⁴ However, encouraging research suggests that relapse, or the recurrence of full-blown symptoms after remission, is rare. In one study, 6 percent of people with borderline personality disorder had a relapse after remission.⁴

Medications :

No medications have been approved by the U.S. Food and Drug Administration to treat borderline personality disorder. Only a few studies show that medications are necessary or effective for people with this illness.³⁵ However, many people with borderline personality disorder are treated with medications in addition to psychotherapy. While medications do not cure BPD, some medications may be helpful in managing specific symptoms. For some people, medications can help reduce symptoms such as anxiety, depression, or aggression. Often, people are treated with several medications at the same time,¹² but there is little evidence that this practice is necessary or effective. Medications can cause different side effects in different people. People who have borderline personality disorder should talk with their prescribing doctor about what to expect from a particular medication.

Other Treatments :

Omega-3 fatty acids. One study done on 30 women with borderline personality disorder showed that omega-3 fatty acids may help reduce symptoms of aggression and depression.³⁶ The treatment seemed to be as well tolerated as commonly prescribed mood stabilizers and had few side effects. Fewer women who took omega-3 fatty acids dropped out of the study, compared to women who took a placebo (sugar pill).

With proper treatment, many people experience fewer or less severe symptoms. However, many factors affect the amount of time it takes for symptoms to improve, so it is important for people with borderline personality disorder to be patient and to receive appropriate support during treatment.

Conclusion:

Border Personality Disorder have cluster of symptoms which is not totally manage. It takes time and different types of treatment available. On the basis of severity of symptoms one can decide which therapy can be used. Professional training and experience required to diagnose and treatment of Borderline Personality Disorder.

References:

1. Gunderson JG. *A BPD Brief: An Introduction to Borderline Personality Disorder: Diagnosis, Origins, Course, and Treatment.* (ed)^(eds).
2. Lenzenweger MF, Lane MC, Loranger AW, Kessler RC. DSM-IV personality disorders in the National Comorbidity Survey Replication. *Biol Psychiatry.* 2007 Sep 15;62(6):553–64.
3. Paris J, Zweig-Frank H. A 27-year follow-up of patients with borderline personality disorder. *Compr Psychiatry.* 2001 Nov–Dec;42(6):482–7.
4. Zanarini MC, Frankenburg FR, Hennen J, Reich DB, Silk KR. The McLean Study of Adult Development (MSAD): overview and implications of the first six years of prospective follow-up. *J Personal Disord.* 2005 Oct;19(5):505–23.
5. Meyer B, Pilkonis PA, Beevers CG. What's in a (neutral) face? Personality disorders, attachment styles, and the appraisal of ambiguous social cues. *J Pers Disord.* 2004 Aug;18(4):320–36.
6. Hazlett EA, Speiser LJ, Goodman M, Roy M, Carrizal M, Wynn JK, Williams WC, Romero M, Minzenberg MJ, Siever LJ, New AS. Exaggerated affect-modulated startle during unpleasant stimuli in borderline personality disorder. *Biol Psychiatry.* 2007 Aug 1;62(3):250–5.
7. Linehan MM, Comtois KA, Murray AM, Brown MZ, Gallop RJ, Heard HL, Korslund KE, Tutek DA, Reynolds SK, Lindenboim N. Two-year randomized controlled trial and follow-up of dialectical behavior therapy vs therapy by experts for suicidal behaviors and borderline personality disorder. *Arch Gen Psychiatry.* 2006 Jul;63(7):757–66.

8. Kleindienst N, Bohus M, Ludascher P, Limberger MF, Kuenkele K, Ebner-Priemer UW, Chapman AL, Reicherzer M, Stieglitz RD, Schmahl C. Motives for nonsuicidal self-injury among women with borderline personality disorder. *J Nerv Ment Dis.* 2008 Mar;196(3):230–6.
9. Chanen AM, Jackson HJ, McCutcheon LK, Jovev M, Dudgeon P, Yuen HP, Germano D, Nistico H, McDougall E, Weinstein C, Clarkson V, McGorry PD. Early intervention for adolescents with borderline personality disorder using cognitive analytic therapy: randomised controlled trial. *Br J Psychiatry.* 2008 Dec;193(6):477–84.
10. Zerkowicz P, Paris J, Guzder J, Feldman R. Diatheses and stressors in borderline pathology of childhood: the role of neuropsychological risk and trauma. *J Am Acad Child Adolesc Psychiatry.* 2001 Jan;40(1):100–5.
11. Zanarini MC, Frankenburg FR, Khera GS, Bleichmar J. Treatment histories of borderline inpatients. *Compr Psychiatry.* 2001 Mar–Apr;42(2):144–50.
12. Zanarini MC. *Ten-Year Course of Borderline Personality Disorder.* (ed)[^](eds). *Borderline Personality Disorder: Course, Outcomes, Interventions.*
13. Tadic A, Wagner S, Hoch J, Baskaya O, von Cube R, Skaletz C, Lieb K, Dahmen N. Gender differences in axis I and axis II comorbidity in patients with borderline personality disorder. *Psychopathology.* 2009;42(4):257–63.
14. Frankenburg FR, Zanarini MC. Obesity and obesity-related illnesses in borderline patients. *J Personal Disord.* 2006 Feb;20(1):71–80.
15. Sansone RA, Hawkins R. Fibromyalgia, borderline personality, and opioid prescription. *Gen Hosp Psychiatry.* 2004 Sep–Oct;26(5):415–6.
16. Torgersen S, Lygren S, Oien PA, Skre I, Onstad S, Edvardsen J, Tambs K, Kringlen E. A twin study of personality disorders. *Compr Psychiatry.* 2000 Nov–Dec;41(6):416–25.
17. Coolidge FL, Thede LL, Jang KL. Heritability of personality disorders in childhood: a preliminary investigation. *J Pers Disord.* 2001 Feb;15(1):33–40.
18. Lynam DR, Widiger TA. Using the five-factor model to represent the DSM-IV personality disorders: an expert consensus approach. *J Abnorm Psychol.* 2001 Aug;110(3):401–12.
19. Lis E, Greenfield B, Henry M, Guile JM, Dougherty G. Neuroimaging and genetics of borderline personality disorder: a review. *J Psychiatry Neurosci.* 2007 May;32(3):162–73.
20. Ruggero CJ, Zimmerman M, Chelminski I, Young D. Borderline personality disorder and the misdiagnosis of bipolar disorder. *J Psychiatr Res.* 2010 Apr;44(6):405–8.
21. Paris J. The diagnosis of borderline personality disorder: problematic but better than the alternatives. *Ann Clin Psychiatry.* 2005 Jan–Mar;17(1):41–6.
22. *Emotion-Regulating Circuit Weakened in Borderline Personality Disorder.* (ed)[^](eds). <http://www.nimh.nih.gov/science-news/2008/emotion-regulating-circuit-weakened-in-borderline-personality-disorder.shtml>. Accessed on Oct 10, 2008.
23. King-Casas B, Sharp C, Lomax-Bream L, Lohrenz T, Fonagy P, Montague PR. The rupture and repair of cooperation in borderline personality disorder. *Science.* 2008 Aug 8;321(5890):806–10.
24. Kernberg OF, Michels R. Borderline personality disorder. *Am J Psychiatry.* 2009 May;166(5):505–8.
25. Silbersweig D, Clarkin JF, Goldstein M, Kernberg OF, Tiescher O, Levy KN, Brendel G, Pan H, Beutel M, Pavony MT, Epstein J, Lenzenweger MF, Thomas KM, Posner MI, Stern E. Failure of frontolimbic inhibitory function in the context of negative emotion in borderline personality disorder. *Am J Psychiatry.* 2007 Dec;164(12):1832–41.
26. Koenigsberg HW, Siever LJ, Lee H, Pizzarello S, New AS, Goodman M, Cheng H, Flory J, Prohovnik I. Neural correlates of emotion processing in borderline personality disorder. *Psychiatry Res.* 2009 Jun 30;172(3):192–9.
27. Binks CA, Fenton M, McCarthy L, Lee T, Adams CE, Duggan C. Psychological therapies for people with borderline personality disorder. *Cochrane Database Syst Rev.* 2006;(1):CD005652.
28. Stone MH. Management of borderline personality disorder: a review of psychotherapeutic approaches. *World Psychiatry.* 2006 Feb;5(1):15–20.
29. Davidson K, Norrie J, Tyrer P, Gumley A, Tata P, Murray H, Palmer S. The effectiveness of cognitive behavior therapy for borderline personality disorder: results from the borderline personality disorder study of cognitive therapy (BOSCOT) trial. *J Personal Disord.* 2006 Oct;20(5):450–65.
30. McMain S, Pos AE. Advances in psychotherapy of personality disorders: a research update. *Curr Psychiatry Rep.* 2007 Feb;9(1):46–52.
31. Kellogg SH, Young JE. Schema therapy for borderline personality disorder. *J Clin Psychol.* 2006 Apr;62(4):445–58.
32. Blum N, St John D, Pfohl B, Stuart S, McCormick B, Allen J, Arndt S, Black DW. Systems Training for Emotional Predictability and Problem Solving (STEPPS) for outpatients with borderline personality disorder: a randomized controlled trial and 1-year follow-up. *Am J Psychiatry.* 2008 Apr;165(4):468–78.
33. Hoffman PD, Fruzzetti AE. Advances in interventions for families with a relative with a personality disorder diagnosis. *Curr Psychiatry Rep.* 2007 Feb;9(1):68–73.
34. McGlashan TH, Grilo CM, Sanislow CA, Ralevski E, Morey LC, Gunderson JG, Skodol AE, Shea MT, Zanarini MC, Bender D, Stout RL, Yen S, Pagano M. Two-year prevalence and stability of individual DSM-IV criteria for schizotypal, borderline, avoidant, and obsessive-compulsive personality disorders: toward a hybrid model of axis II disorders. *Am J Psychiatry.* 2005 May;162(5):883–9.
35. Binks CA, Fenton M, McCarthy L, Lee T, Adams CE, Duggan C. Pharmacological interventions for people with borderline personality disorder. *Cochrane Database Syst Rev.* 2006;(1):CD005653.
36. Zanarini MC, Frankenburg FR. omega-3 Fatty acid treatment of women with borderline personality disorder: a double-blind, placebo-controlled pilot study. *Am J Psychiatry.* 2003 Jan;160(1):167–9.